



General Certificate of Education

Statistics 6380

SS03 Statistics 3

Mark Scheme

2007 examination - June series

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation meeting attended by all examiners and is the scheme which was used by them in this examination. The standardisation meeting ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for the standardisation meeting each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed at the meeting and legislated for. If, after this meeting, examiners encounter unusual answers which have not been discussed at the meeting they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available to download from the AQA Website: www.aqa.org.uk

Copyright © 2007 AQA and its licensors. All rights reserved.

COPYRIGHT

AQA retains the copyright on all its publications. However, registered centres for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to centres to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

Key to mark scheme and abbreviations used in marking

M	mark is for method		
m or dM	mark is dependent on one or more M marks and is for method		
A	mark is dependent on M or m marks and is for accuracy		
B	mark is independent of M or m marks and is for method and accuracy		
E	mark is for explanation		
✓ or ft or F	follow through from previous incorrect result	MC	mis-copy
CAO	correct answer only	MR	mis-read
CSO	correct solution only	RA	required accuracy
AWFW	anything which falls within	FW	further work
AWRT	anything which rounds to	ISW	ignore subsequent work
ACF	any correct form	FIW	from incorrect work
AG	answer given	BOD	given benefit of doubt
SC	special case	WR	work replaced by candidate
OE	or equivalent	FB	formulae book
A2,1	2 or 1 (or 0) accuracy marks	NOS	not on scheme
-x EE	deduct x marks for each error	G	graph
NMS	no method shown	c	candidate
PI	possibly implied	sf	significant figure(s)
SCA	substantially correct approach	dp	decimal place(s)

No Method Shown

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded. However, there are situations in some units where part marks would be appropriate, particularly when similar techniques are involved. Your Principal Examiner will alert you to these and details will be provided on the mark scheme.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

Otherwise we require evidence of a correct method for any marks to be awarded.

SS03

Q	Solution	Marks	Total	Comments																						
1	<p>H_0 (pop) median/mean diff $\eta_d = 0$ H_1 (pop) median/mean diff $\eta_d \neq 0$ 2 tail 5%</p> <table border="1"> <tr> <td>Difference (without – with)</td> <td>4</td> <td>2</td> <td>-5</td> <td>0</td> <td>4</td> </tr> <tr> <td>Rank</td> <td>4½</td> <td>2</td> <td>-6</td> <td>.</td> <td>4½</td> </tr> </table> <table border="1"> <tr> <td>-6</td> <td>-3</td> <td>-7</td> <td>-9</td> <td>1</td> </tr> <tr> <td>-7</td> <td>-3</td> <td>-8</td> <td>-9</td> <td>1</td> </tr> </table> <p>$T_+ = 4\frac{1}{2} + 2 + 4\frac{1}{2} + 1 = 12$ $T_- = 6 + 7 + \dots + 9 = 33$</p> <p>test statistic $T = 12$ $cv = 6$ $n = 9$ $T > 6$</p> <p>No significant evidence at 5% level to reject H_0. Accept H_0 There is no significant evidence to suggest that customers rated differently the radio after introducing the new component</p>	Difference (without – with)	4	2	-5	0	4	Rank	4½	2	-6	.	4½	-6	-3	-7	-9	1	-7	-3	-8	-9	1	<p>B1</p> <p>M1</p> <p>M1</p> <p>m1</p> <p>A1 B1 M1✓</p> <p>A1</p> <p>E1</p>	<p>9</p>	<p>Need ‘average’</p> <p>For differences (+/- signs can be interchanged); ignore signs</p> <p>For ranks. Rank 1=1</p> <p>For totals</p> <p>For one correct total For $cv = 6$ Comparison cv/ts</p> <p>In context (ft)</p>
Difference (without – with)	4	2	-5	0	4																					
Rank	4½	2	-6	.	4½																					
-6	-3	-7	-9	1																						
-7	-3	-8	-9	1																						
	Total		9																							
2(a)	<p>From calculator $r = 0.915$ (0.91456)</p> $\text{or } r = \frac{2102.57 - \left(\frac{135.2 \times 147.9}{10}\right)}{\sqrt{128.976} \times \sqrt{98.269}}$ $= \frac{102.962}{11.35 \times 9.913}$ $= 0.915$	B3	3	<p>AWRT B2 for 0.914 or 0.91 – 0.92 B1 for 0.9</p> <p>Alternative: $n = 10$ $\sum x = 135.2$ $\sum y = 147.9$ $\sum x^2 = 1956.88$ $\sum y^2 = 2285.71$ $\sum xy = 2102.57$ (M1) sub in formula (m1) (A1)</p>																						
(b)	<p>$H_0 \rho = 0$ $H_1 \rho > 0$ 1 tail 1% sig level</p> <p>test statistic $r = 0.915$ $cv = 0.7155$ $n = 10$ since $ts > 0.7155$</p> <p>Reject H_0</p> <p>Significant evidence at 1% level to suggest a positive linear association between the weight gain of mothers during pregnancy and the weight of their children at 3 years of age</p>	<p>B1</p> <p>B1 M1 A1</p> <p>E1</p>	<p>5</p>	<p>Or words</p> <p>For cv For comparison ts/cv</p> <p>In context (ft)</p>																						
	Total		8																							

SS03 (cont)

Q	Solution	Marks	Total	Comments
3(a)(i)	H ₀ No association between height at one year old and income at age 50 years H ₁ An association exists between height at one year old and income at age 50 years 1 tail 5%	B1		H ₀ independent H ₁ not independent
3(a)(ii)	Those babies with a low height, under 75 cm, at age one year appear more likely to achieve a lower income at age 50 years and those babies with heights 80 cm and over at age one year appear more likely to achieve an income of £20,500 and over	E1		Must have attempted χ^2
		E1	2	Indication of sources of association in context

	Under £20,500	£20,500 and over
Under 75 cm	8.5	11.5
75 cm to under 80 cm	12.75	17.25
80 cm and over	12.75	17.25

$$ts = \sum \frac{(O - E)^2}{E} = \frac{5.5^2}{8.5} + \frac{5.5^2}{11.5} + \frac{0.75^2}{12.75} + \frac{0.75^2}{17.25} + \frac{4.75^2}{12.75} + \frac{4.75^2}{17.25}$$

$$= 3.56 + 2.63 + 0.044 + 0.033 + 1.77 + 1.31 = 9.34$$

df = 2 5% cv = 5.991
ts > 5.991

Reject H₀
Significant evidence to suggest an association exists between height at one year old and income at age 50 years

M1
m1

M1
A1
B1
m1✓

A1

8

E method for 3 correct
For all E correct

ts sum with correct denominators

For ts in range 9.10 – 9.50
For cv
For comparison ts/cv

For **reject H₀**

SS03 (cont)

Q	Solution	Marks	Total	Comments									
3(b)	H ₀ Presence of parent is independent of occurrence of emergency medical treatment H ₁ Presence of parent is not independent of occurrence of emergency medical treatment 1 tail 5%	B1		Or as in (a)									
	<table border="1"> <thead> <tr> <th></th> <th>Travelled</th> <th>Did not travel</th> </tr> </thead> <tbody> <tr> <th>Required</th> <td>12.490</td> <td>5.510</td> </tr> <tr> <th>Not required</th> <td>89.510</td> <td>39.490</td> </tr> </tbody> </table>		Travelled	Did not travel	Required	12.490	5.510	Not required	89.510	39.490	M1		For E values method
		Travelled	Did not travel										
	Required	12.490	5.510										
	Not required	89.510	39.490										
	$ts = \sum \frac{(O - E - 0.5)^2}{E}$	M1		For ts									
	$= \frac{1.990^2}{12.490} + \frac{1.990^2}{5.510} + \frac{1.990^2}{89.510} + \frac{1.990^2}{39.490}$	m1		For Yates' corr									
$= 0.317 + 0.719 + 0.044 + 0.100$	A1		For ts 1.00 – 1.30										
$= 1.18$ df = 1 5% cv = 3.84 ts < 3.84 Accept H ₀ No significant evidence to doubt that presence of parent is independent of occurrence of emergency medical treatment	B1 M1✓		For cv For comparison ts/cv										
	A1	8	In context										
	Total		18										

SS03 (cont)

Q	Solution	Marks	Total	Comments																			
4(a)	H_0 (pop) median $\eta = 14$ H_1 (pop) median $\eta > 14$	B1		Not mean																			
	2 tail 10% Signs: - + + + + + + - + $n = 9$ test stat = $7^+ / 2^-$	M1		Signs SC2: Wilcoxon signed-rank																			
	Model B(9, 0.5)	A1		test stat correct																			
	$P(\leq 2^-) = P(\geq 7^+) = 0.0898 < 0.10$	M1		Bin model seen to be used ($n = 9, p = 0.5$ column)																			
	Reject H_0 Significant evidence at 10% level to doubt H_0 There is significant evidence to suggest that the median cocaine use has increased since 2000	M1		Comparison of correct B(9, 0.5) probability with 0.05 or 0.10 Or use of identified cv cr [7, 8, 9] see 0.0898																			
		A1																					
		E1	7																				
	(b)(i)	<table border="1"> <thead> <tr> <th>Town</th> <th>A</th> <th>B</th> <th>C</th> <th>D</th> <th>E</th> </tr> </thead> <tbody> <tr> <td>x rank</td> <td>9</td> <td>1</td> <td>5</td> <td>4</td> <td>2</td> </tr> <tr> <td>y rank</td> <td>1</td> <td>8</td> <td>5</td> <td>6</td> <td>9</td> </tr> </tbody> </table>	Town	A	B	C	D	E	x rank	9	1	5	4	2	y rank	1	8	5	6	9	M1		Attempt at ranks
		Town	A	B	C	D	E																
		x rank	9	1	5	4	2																
y rank		1	8	5	6	9																	
<table border="1"> <thead> <tr> <th>Town</th> <th>F</th> <th>G</th> <th>H</th> <th>I</th> <th></th> </tr> </thead> <tbody> <tr> <td>x rank</td> <td>6</td> <td>3</td> <td>8</td> <td>7</td> <td></td> </tr> <tr> <td>y rank</td> <td>3</td> <td>7</td> <td>2</td> <td>4</td> <td></td> </tr> </tbody> </table>	Town	F	G	H	I		x rank	6	3	8	7		y rank	3	7	2	4		M1		x or y ranks correct (reverse order OK)		
Town	F	G	H	I																			
x rank	6	3	8	7																			
y rank	3	7	2	4																			
	A1			All correct, consistent Rank all as one M1M1 only																			
	$r_s = -0.967$ (3 sf from calc) (-0.966 to -0.967 B3) (-0.96 to -0.97 B2)	B3	6	Alternative: $d = 8, 7, 0, 2, 7, 3, 4, 6, 3$ $\sum d^2 = 236$ M1 $r_s = 1 - \frac{6 \times 236}{9 \times 80} = -0.967$ m1A1																			
(ii)	The estimated cocaine use in a town is higher when it is easier to buy cocaine in the town	E1	1	Must have some sensible answer in (i); comparative required																			
	Total		14																				

SS03 (cont)

Q	Solution	Marks	Total	Comments																						
5(a)	H ₀ Samples are taken from identical populations H ₁ Samples are not taken from identical populations – population average scores differ	B1		Hypotheses referring to population averages also acceptable or fully explained in words																						
	<table border="1"> <thead> <tr> <th>Morning</th> <th>Afternoon</th> </tr> </thead> <tbody> <tr><td>44</td><td>46</td></tr> <tr><td>53</td><td>47</td></tr> <tr><td>54</td><td>51</td></tr> <tr><td>56</td><td>58</td></tr> <tr><td>63</td><td>59</td></tr> <tr><td>63</td><td>61</td></tr> <tr><td>65</td><td>62</td></tr> <tr><td>72</td><td>67</td></tr> <tr><td>74</td><td>68</td></tr> <tr><td>81</td><td></td></tr> </tbody> </table>	Morning	Afternoon	44	46	53	47	54	51	56	58	63	59	63	61	65	62	72	67	74	68	81		M1		Separation of am/pm
	Morning	Afternoon																								
	44	46																								
	53	47																								
	54	51																								
	56	58																								
	63	59																								
	63	61																								
	65	62																								
72	67																									
74	68																									
81																										
Ranks: Afternoon 2 3 4 8 9 10 11 15 16 Morning 1 5 6 7 12½ 12½ 14 17 18 19	M1 A1		Or reversed																							
T _A = 2 + 3 + + 16 = 78 T _M = 1 + 5 + + 19 = 112	m1ft		102 88 Or alt method directly to U																							
$U_A = 78 - \frac{9 \times 10}{2} = 33$	m1																									
$U_M = 112 - \frac{10 \times 11}{2} = 57$																										
Test stat U = 33 cv = 21 n = 9, m = 10 U = 33 > 21	A1 B1 M1		U = 33 or U = 57 Comparison U/cv; not if U < 0																							
Accept H ₀ No significant evidence at the 5% level to suggest that there is any difference in average test scores between students taking the test in a morning or afternoon session	E1	10	In context																							
(b)(i) In matched pairs design, individual differences are minimised since the same person is tested each time and therefore any difference which may exist between the two groups is more likely to be identified	B1 B1	2	Reduce experimental error; avoid bias																							
(ii) She kept the students apart during the day of the test She chose students of similar ability and randomly assigned them to a morning or an afternoon session	E1 E1	2	Any 2																							
	Total		14																							

SS03 (cont)

Q	Solution	Marks	Total	Comments																					
6	<p>H_0 Samples are taken from identical populations H_1 Samples are not taken from identical populations – population average protective antibody levels differ</p> <p>Ranks:</p> <table border="1"> <thead> <tr> <th>Happily Married</th> <th>Unhappily Married</th> <th>Unmarried</th> </tr> </thead> <tbody> <tr> <td>7</td> <td>5</td> <td>1</td> </tr> <tr> <td>10</td> <td>6</td> <td>2</td> </tr> <tr> <td>12</td> <td>9</td> <td>3</td> </tr> <tr> <td>13</td> <td>11</td> <td>4</td> </tr> <tr> <td>14</td> <td></td> <td>8</td> </tr> <tr> <td>15</td> <td></td> <td></td> </tr> </tbody> </table> <p> $T_{Hap} = 71$ $T_{Unhap} = 31$ $T_{Unmarr} = 18$ $n_{Hap} = 6$ $n_{Unhap} = 4$ $n_{Unmarr} = 5$ </p> $\sum_{i=1}^m \frac{T_i^2}{n_i} = \frac{71^2}{6} + \frac{31^2}{4} + \frac{18^2}{5} = 1145.22$ $H = \frac{12}{15 \times 16} \times 1145.22 - (3 \times 16)$ <p>= 9.26</p> <p>Critical value from $\chi^2_2 = 5.991$ $H > 5.991$</p> <p>Sig evidence to reject H_0 and conclude that samples are not from identical populations</p> <p>Significant evidence at the 5% level to suggest that the population average level of protective antibodies differs for the three marital categories: at least two of the averages differ</p> <p>It appears that those males who are happily married have a significantly higher level of antibodies than those who are unmarried</p>	Happily Married	Unhappily Married	Unmarried	7	5	1	10	6	2	12	9	3	13	11	4	14		8	15			<p>B1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>m1</p> <p>m1</p> <p>A1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>E1</p> <p>E1</p>	<p>12</p>	<p>or $H_0 \eta_{Happy} = \eta_{Unhappy} = \eta_{Unmarried}$ H_1 at least two of $\eta_{Happy}, \eta_{Unhappy}, \eta_{Unmarried}$ do differ</p> <p>Allow mean B1 H_0 antibody independent of marital status H_1 antibody not independent of marital status</p> <p>For 10</p> <p>Totals</p> <p>Correct method test stat:</p> $H = \frac{12}{N(N+1)} \sum_{i=1}^m \frac{T_i^2}{n_i} - 3(N+1)$ <p>9.1–9.4</p> <p>Difference in context Mention of ‘at least two’ or happily married/unmarried differ</p>
Happily Married	Unhappily Married	Unmarried																							
7	5	1																							
10	6	2																							
12	9	3																							
13	11	4																							
14		8																							
15																									
	Total		12																						
	TOTAL		75																						